



## Why Tropical Trees Have Rotten Cores

Daniel H. Janzen

*Biotropica*, Volume 8, Issue 2 (Jun., 1976), 110.

---

Your use of the JSTOR database indicates your acceptance of JSTOR's Terms and Conditions of Use. A copy of JSTOR's Terms and Conditions of Use is available at <http://www.jstor.org/about/terms.html>, by contacting JSTOR at [jstor-info@umich.edu](mailto:jstor-info@umich.edu), or by calling JSTOR at (888)388-3574, (734)998-9101 or (FAX) (734)998-9113. No part of a JSTOR transmission may be copied, downloaded, stored, further transmitted, transferred, distributed, altered, or otherwise used, in any form or by any means, except: (1) one stored electronic and one paper copy of any article solely for your personal, non-commercial use, or (2) with prior written permission of JSTOR and the publisher of the article or other text.

Each copy of any part of a JSTOR transmission must contain the same copyright notice that appears on the screen or printed page of such transmission.

*Biotropica* is published by Association for Tropical Biology. Please contact the publisher for further permissions regarding the use of this work. Publisher contact information may be obtained at <http://www.jstor.org/journals/tropbio.html>.

---

*Biotropica*

©1976 Association for Tropical Biology

JSTOR and the JSTOR logo are trademarks of JSTOR, and are Registered in the U.S. Patent and Trademark Office. For more information on JSTOR contact [jstor-info@umich.edu](mailto:jstor-info@umich.edu).

©2001 JSTOR

- A. Allen, H. Lang, and J. P. Chapin, The American museum Congo expedition collection. Bull. Am. Mus. nat. Hist. 37: 479-563.
- LEIN, M. R. 1972. A trophic comparison of avifaunas. Syst. Zool. 21: 135-150.
- MACDONALD, D. 1973. Birds of Australia, a summary of information. A. H. and A. W. Reed, Sydney.
- MACKWORTH-PRAED, C. W. 1970. Birds of west and central and western Africa. African handbook of birds Series III, Volume 1. Longman, London.
- MARTI, C. D. 1974. Feeding ecology of four sympatric owls. Condor 76: 45-61.
- MEDWAY, LORD. 1969. The wild mammals of Malaya and offshore islands, including Singapore. Oxford Univ. Press, London.
- NOVICK, A., AND B. A. DALE. 1971. Foraging behavior in fishing bats and their insectivorous relatives. J. Mammal. 52: 817-818.
- PETERS, J. L. 1940. Check-list of birds of the world. Volume IV. Harvard Univ. Press, Cambridge.
- PRAKASH, I. 1959. Foods of the Indian false vampire. J. Mammal. 40: 545-547.
- RIDE, W. D. L. 1970. A guide to the native mammals of Australia. Oxford Univ. Press, London.
- ROSEVEAR, D. R. 1965. The bats of West Africa. Brit. Mus. Nat. Hist. London.
- ROSS, A. 1969. Ecological aspects of the food habits of insectivorous screech-owls. Proc. Western Found. Vert. Zool. 1: 301-344.
- ROTHSCHILD, M. 1971. Speculations about mimicry with Henry Ford. In, R. Creed (Ed.), Ecological genetics and evolution, pp. 202-223. Blackwell Sci. Pub., Oxford.
- SWYNNERTON, C. F. 1915. Birds in relation to their prey: experiments on wood-hoopoes, small hornbills and a babler. J. S. Afr. Orn. Un. 11: 32-108.
- VAN DEUSEN, H. M. 1968. Carnivorous habits of *Hypsignathus monstrosus*. J. Mammal. 49: 335-336.
- VAN GELDER, R. G., AND W. W. GOODPASTER. 1952. Bats and birds competing for food. J. Mammal. 33: 491.
- VAN TYNE, J., AND A. J. BERGER. 1959. Fundamentals of Ornithology. John Wiley and Sons, New York.
- WALKER, E. P. 1964. Mammals of the World, Volume 1. Johns Hopkins Press, Baltimore.
- WILSON, D. E. 1973. Bat faunas: a trophic comparison. Syst. Zool. 22: 14-29.
- 

## NOTE

### Why Tropical Trees Have Rotten Cores

It is generally assumed that a rotten core in an adult tree is detrimental. On the contrary, I hypothesize that the rotten hollow core is often an adaptive trait, selected for as a mechanism of nitrogen and mineral trapping. A rotten core is a site of animal nests, animal defecation, and microbial metabolism that should result in a steady fertilization of the soil under the base of the tree. If it can be shown that none of this natural fertilizer is harvested by the tree's own roots, then the hypothesis is rejected. Theoretically, a tree could be programmed for a rotten core through the mechanism of depositing amounts and kinds of heartwood secondary compounds that would be sufficient to exclude organisms until the tree is large enough to tolerate structurally a rotten core, with subsequent access by rotting organisms and others to the central part of this core through unhealed breaks in branches, roots, or trunk. A complementary trait should be an increased deposition in the outer heartwood, a deposition that would contain the rotten-core organisms and therefore protect the sapwood and structural soundness of the tree. Such a hypothesis assumes that as a tree reaches adult size, it need not grow an ever-larger solid trunk and that a hollow core is not structurally lethal. Hollow cores are expected most frequently in nutrient-poor sites, habitats, or life forms not plagued by high winds, and in situations containing animals and microorganisms that will use a hollow core without overwhelming the living tree. In short, the hollow core becomes a clever use of an otherwise useless piece of wood.

**Daniel H. Janzen**

Department of Biology  
University of Pennsylvania  
Philadelphia, Pennsylvania 19174 U.S.A.

---